

MATH203 Calculus

Dr. Bandar Al-Mohsin

School of Mathematics, KSU

Area and Volume

Volume:

In the previous study, we saw that if $f(x, y) \geq 0$ and f is continuous, then the double integral

$$\iint_R f(x, y) dA \quad (1)$$

gives the volume of the solid that lies under the graph of $z = f(x, y)$ and over a region R in the xy -plane.

Area:

The double integral (1) can be used to find the area of the region R if $f(x, y) = 1$ which becomes


$$\iint_R dA \quad (2)$$

Area and Volume

Double Integral for finding area:

Formula 1 If a region R_x is defined by $a \leq x \leq b$ and $g(x) \leq y \leq h(x)$, where $g(x), h(x)$ are continuous on $[a, b]$, then the area A of R_x is given by


$$A = \int_a^b \int_{g(x)}^{h(x)} dy dx$$



Area and Volume

Formula 2 If a region R_y is defined by $c \leq y \leq d$ and $p(y) \leq x \leq q(y)$, where $p(y), q(y)$ are continuous on $[c, d]$, then the area A of R_y is given by

$$\int_c^d \int_{p(y)}^{q(y)} dx dy$$



Double Integrals

Examples

Sketch the region bounded by the graphs of :

(1) $y = x^2$ and $y = 2x$. Evaluate $\iint_R (x^3 + 4y) dA$ using R_x region and

R_y region.

(2) $y = \sqrt{x}$ and $y = \sqrt{3x - 18}$ and $y = 0$ using R_x region and R_y region.

(3) reverse the order of the integration and evaluate

$$\int_0^4 \int_{\sqrt{y}}^2 y \cos x^5 dx dy.$$

(4) Find the area A of the region in the xy -plane bounded by the graph of $x = y^3$, $x + y = 2$ and $y = 0$

(4) Find the volume of the solid that lies under the graph of $z = 4x^2 + y^2$ and over the region in the xy -plane bounded by $x + y = 2$, $x = 0$, $y = 0$ and $z = 0$.